

References

- MARZOLF, E. R., P. J. MULHOLLAND, AND A. D. STEINMAN. 1994. Improvements to the diurnal upstream-downstream dissolved oxygen change technique for determining whole-stream metabolism in small streams. *Can. J. Fish. Aquat. Sci.* **51**: 1591–1599.
- , ———, AND ———. 1998. Reply: Improvements to the diurnal upstream-downstream dissolved oxygen change technique for determining whole-stream metabolism in small streams. *Can. J. Fish. Aquat. Sci.* **55**: 1786–1787.
- MULHOLLAND, P. J., E. R. MARZOLF, J. R. WEBSTER, D. R. HART, AND S. P. HENDRICKS. 1997. Evidence that hyporheic zones increase heterotrophic metabolism and phosphorus uptake in forest streams. *Limnol. Oceanogr.* **42**: 443–451.
- YOUNG, R. G., AND A. D. HURYN. 1998. Comment: Improvements to the diurnal upstream-downstream dissolved oxygen change technique for determining whole-stream metabolism in small streams. *Can. J. Fish. Aquat. Sci.* **55**: 1784–1785.

Received: 13 October 1998

Size-dependent C:N uptake by phytoplankton as a function of irradiance: Ecological implications

Our paper published in *Limnology and Oceanography* (Vol. 43, p. 1362–1368) contains a significant typographic error. On page 1365, column 2, line 7 onwards should read: “For the south basin, only the POC:PON data for the <2- μm fraction was significantly correlated with $1/k$ ($r = -0.547$, $P = 0.0284$, $n = 16$). There was no overall correlation between the POC:PON ratio for the >2- μm fraction

and transparency in the south basin ($r = +0.218$, $P = 0.434$, $n = 15$), . . .”.

J.-J. Frenette, W. F. Vincent, and L. Legendre

Received: 12 November 1998